2 Which equation represents a parabola with a focus of (0,4) and a directrix of $y = 2$?

(1) $y \quad x^2 \quad 3$

(2) $y \quad x^2 \quad 1$

(3) $y \quad \frac{x^2}{2} \quad 3$

(4) $y \quad \frac{x^2}{4} \quad 3$

3 If the terminal side of angle , in standard position, passes through point (–4,3), what is the numerical value of sin ?

(1) $\frac{3}{5}$

(2) $\frac{4}{5}$

(3) $-\frac{3}{5}$

(4) $-\frac{4}{5}$

5 Use the properties of rational exponents to determine the value of *y* for the equation:

$$\frac{\sqrt[3]{x^8}}{x^{\frac{1}{3}}} \quad x^y,\ x \neq 1.$$

Key: $\frac{4}{3}$

Measures CCLS Cluster: N-RN.A

Mathematical Practice: 2

Commentary: This question measures N-RN.A because students extend their knowledge of integer exponents to rewriting and working with radicals in terms of rational exponents.

Rationale:

$$\frac{x^{\frac{8}{3}}}{x^{\frac{4}{3}}} \quad x^y$$

$$x^{\frac{4}{3}} \quad x^y$$

$$\frac{4}{3} \quad y$$

Rubric:

[2] $\frac{4}{3}$ or an equivalent value is written, and correct work using rational exponents is shown.

[1] Appropriate work is shown, but one computational error is made.

or

[1] Appropriate work is shown, but one conceptual error is made.

or

[1] $\frac{x^{\frac{8}{3}}}{x^{\frac{4}{3}}}$ is written, but no further correct work is shown.

or

[1] $\frac{4}{3}$, but no work is shown.

[0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.

6 Write $5 \quad 2yi \quad 4 \quad 3i \quad\quad 5 \quad 2yi \quad 4 \quad 3i$ in

8 Solve algebraically for all values of x:

$$\sqrt{x \quad 5} \quad x \quad 7$$

Key: 6

Measures CCLS Cluster: A-REI.A

Mathematical Practice: 3, 6

Commentary: This question measures A-REI.A because the problem requires students to solve a radical equation and identify extraneous solutions.

Rationale:

$$x$$

$$7 \quad x^{2}$$

$$x \quad 5 \quad 49 \quad 14x \quad x^{2}$$

$$x^{2} \quad o2\ 1\ 6$$

11 Write an explicit formula for a_n, the nth term of the recursively defined sequence below.

$$a_1 \quad x \quad 1$$
$$a_n \quad x \quad a_{n\ 1}$$

For what values of x would $a_n \quad 0$ when $n\ !\ 1$?